K-Max Lite 0-0-14.5

Guaranteed Analysis:
Soluble Potash (K_2O) 14.5%
Derived from potassium chloride.

A Potassium True Solution for use in Organic Production

For use in certified organic production as a fertilizer in foliar, fertigation and soil applications to supplement potassium levels on agricultural crops.
Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85ºF before application completely dries on foliage.
- Aerial application and ground sprays – minimum of 5 gallons of water/acre; Dilute sprays – minimum of 30 gallons of water/acre.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- K-Max Lite is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- Care should be exercised when applying to chlorophobic plants.
- **Consult a NutriAg representative for application timing and necessity of its use.**

Suggested Application Rates:

Crop	Foliar Rates**	Fertigation Rates**
Forage Crops: Alfalfa, Clover, Hay Crops **Field Crops:** Cereals, Corn, Cotton, Sorghum, Sugar cane **Legumes:** Dry bean, Lentil, Peanut, Pea, Soybean **Root Vegetables (including, but not limited to):** Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip **Leafy Vegetables (including but not limited to):** Bok Choy, Celery, Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach **Other Vegetables:** Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato **Berry Crops:** Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry **Tree Fruit (including but not limited to):** Apple, Apricot, Avocado, Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince **Other Fruit:** Citrus (all), Fig, Kiwi, Melons **Nut Trees (including but not limited to):** Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut **Grapes:** Currant, Raisin, Table, Wine **Other Crops:** Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower	1 – 4 quarts/acre/ application	The number and frequency of applications determine application rates. Apply no more than 2 gallons/acre/ application. Always apply toward the end of an irrigation cycle.

****Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.**

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg Ltd. warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg Ltd. NutriAg Ltd. will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

The use of the fertilizing material on any crops other than those recommended may result in serious injury to crop(s).

Caution: KEEP OUT OF REACH OF CHILDREN.

OMRI Listed

Listed by the Organic Materials Review Institute (OMRI) for use in organic production

REGISTERED cdfa ORGANIC INPUT MATERIAL

Net Contents:
2.5 Gal. (9.46 L)

Net Weight:
24.60 lbs. (11.15 kg)

Net Weight/Gallon:
9.84 lbs. (4.46 kg)

Batch No.: XXXXXX

SPECIMEN